arXiv:1003.3729v1 [math.QA] 19 Mar 2010

THE QUANTUM CASIMIR OPERATORS OF $\mathrm{U}_q(\mathfrak{gl}_n)$ AND THEIR EIGENVALUES

JUNBO LI

ABSTRACT. We show that the quantum Casimir operators of the quantum linear group constructed in early work of Bracken, Gould and Zhang together with one extra central element generate the entire center of $\mathrm{U}_q(\mathfrak{gl}_n)$. As a by product of the proof, we obtain intriguing new formulae for eigenvalues of these quantum Casimir operators, which are expressed in terms of the characters of a class of finite dimensional irreducible representations of the classical general linear algebra.

Keywords: Quantum groups, Harish-Chandra isomorphism, Center

MR(2010) Subject Classification: 17B10, 17B37

1. INTRODUCTION

Quantum groups originated from the theory of soluble models of the Yang-Baxter type in the middle of 80s. They have played important roles in various branches of mathematics and physics, most notably in two-dimensional soluble models in statistical mechanics and knot theory. The study of their structure and representation theory has been the focus of research in representation theory and continues to attract much attention. In particular, the discovery of the crystal basis and canonical basis [3, 5] is one of the most important achievement in representation theory in recent years.

In the early 90s, a set of central elements of the quantum group $\mathrm{U}_q(\mathfrak{gl}_n)$ was constructed in [8, 9]. We shall refer to them as the quantum Casimir operators of $\mathrm{U}_q(\mathfrak{gl}_n)$. The reason for this terminology is the fact that these central elements of $\mathrm{U}_q(\mathfrak{gl}_n)$ are the quantum analogues of the familiar Casimir operators of the universal enveloping algebra $\mathrm{U}(\mathfrak{gl}_\mathfrak{n})$ of $\mathfrak{gl}_\mathfrak{n}$ given by $\sum_{i_1=1}^{n}\cdots\sum_{i_k=1}^{n}E_{i_1i_2}E_{i_2i_3}\cdots E_{i_{k-1}i_k}E_{i_ki_1}$ $(k=1,2,\cdots)$, where E_{ij} are the images of the matrix units under the canonical embedding of $\mathfrak{gl}_\mathfrak{n}$ in $\mathrm{U}(\mathfrak{gl}_\mathfrak{n})$. These Casimir operators of $\mathrm{U}(\mathfrak{gl}_\mathfrak{n})$ play an important role in the Interacting Boson Model in nuclear physics. Their quantum analogues have also been applied in a similar way.

One obvious question was whether the quantum Casimir operators [8, 9] of $\mathrm{U}_q(\mathfrak{gl}_n)$ (supplemented with the obvious central element c given by (2.2)) generated the entire center of $\mathrm{U}_q(\mathfrak{gl}_n)$. The general expectation was that the answer should be affirmative, but no proof was ever given as far as we know. The main purpose of this paper is to give a rigorous proof. The result is described in Theorem 4.1.

The proof of Theorem 4.1 requires us to analyze the eigenvalues of the quantum Casimir operators in highest weight representations of $\mathrm{U}_q(\mathfrak{gl}_n)$. A formula for the eigenvalues was obtained in [4] (in fact [4] treated $\mathrm{U}_q(\mathfrak{gl}_{\mathfrak{m}|\mathfrak{n}})$, which included $\mathrm{U}_q(\mathfrak{gl}_n)$ as a special case). We cast the formula into a form readily usable for our purpose. This new formula is expressed in terms of the characters of a class of finite dimensional irreducible representations of the classical general linear algebra. This result is rather intriguing, and we believe that it is interesting in its own right.

We should point out that the structure of the center of a quantum group is much studied [2, 7] at an abstract level. In particular, a quantum analogue of the celebrated Harish-Chandra homomorphism in semi-simple Lie algebras has been established for quantum groups at generic q. In the case of $\mathrm{U}_q(\mathfrak{gl}_n)$, a set of generators different from the quantum Casimir operators of [8] was constructed in [1].

2. THE QUANTUM GENERAL LINEAR GROUP

2.1. **The quantum general linear group.**

This section provides some basic materials on the general linear algebra $\mathfrak{gl}_n$ and its quantum group $\mathrm{U}_q(\mathfrak{gl}_n)$. Let ε_i, with $i \in \mathbf{I} = \{1,2,\cdots,n\}$, be a basis of an Euclidean space with the inner product $(\varepsilon_i,\varepsilon_j) = \delta_{ij}$. Set $\rho = \frac{1}{2}\sum_{i=1}^{n}(n-2i+1)\varepsilon_i$.

The quantized universal enveloping algebra $\mathrm{U}_q(\mathfrak{gl}_n)$ of the general linear Lie algebra $\mathfrak{gl}_n$ is a unital associative algebra over $\mathbb{K} := \mathbb{C}(q)$ generated by $K_i^{\pm 1}$, $E_{i'}$, $F_{i'}$ $(i \in \mathbf{I}, i' \in \mathbf{I}' := \mathbf{I}\backslash\{n\})$, subject to the following relations:

$$\begin{aligned}
&K_iK_i^{-1} = 1, \quad K_i^{\pm 1}K_j^{\pm 1} = K_j^{\pm 1}K_i^{\pm 1}, \quad K_iE_jK_i^{-1} = q^{(\varepsilon_j-\varepsilon_{j+1},\varepsilon_i)}E_j,\\
&K_iF_jK_i^{-1} = q^{-(\varepsilon_j-\varepsilon_{j+1},\varepsilon_i)}F_j, \quad E_rE_s = E_sE_r, \ \ F_rF_s = F_sF_r, \quad |r-s| \geqslant 2,\\
&E_iF_j - F_jE_i = \delta_{i,j}\frac{K_iK_{i+1}^{-1} - K_i^{-1}K_{i+1}}{q-q^{-1}}, \quad S_{i,i\pm 1}^{(+)} = S_{i,i\pm 1}^{(-)} = 0,
\end{aligned}$$

and

$$\begin{aligned}
S_{i,i\pm 1}^{(+)} &= (E_i)^2E_{i\pm 1} - (q+q^{-1})E_iE_{i\pm 1}E_i + E_{i\pm 1}(E_i)^2,\\
S_{i,i\pm 1}^{(-)} &= (F_i)^2F_{i\pm 1} - (q+q^{-1})F_iF_{i\pm 1}F_i + F_{i\pm 1}(F_i)^2.
\end{aligned}$$

As we know, $\mathrm{U}_q(\mathfrak{gl}_n)$ possesses the structure of a Hopf algebra with the co-multiplication Δ, co-unit ε and antipode S respectively given by

$$\begin{aligned}
&\Delta(E_i) = E_i \otimes K_iK_{i+1}^{-1} + 1 \otimes E_i, \quad \Delta(F_i) = F_i \otimes 1 + K_i^{-1}K_{i+1} \otimes F_i,\\
&\Delta(K_i^{\pm 1}) = K_i^{\pm 1} \otimes K_i^{\pm 1}, \quad \varepsilon(E_i) = \varepsilon(F_i) = 0, \ \ \varepsilon(K_i^{\pm 1}) = 1,\\
&S(E_i) = -E_iK_i^{-1}K_{i+1}, \quad S(F_i) = -K_iK_{i+1}^{-1}F_i, \quad S(K_i^{\pm 1}) = K_i^{\mp 1}.
\end{aligned}$$

The natural module V for $\mathrm{U}_q(\mathfrak{gl}_n)$ has the standard basis $\{v_i \,|\, i \in \mathbf{I}\}$ such that

$$E_iv_j = \delta_{j,i+1}v_i, \quad F_iv_j = \delta_{j,i}v_{i+1}, \quad K_iv_j = \big(1+(q-1)\delta_{i,j}\big)v_j.$$

Denote π the $\mathrm{U}_q(\mathfrak{gl}_n)$-representation relative to this basis, then $\pi(E_i) = E_{i,i+1}$, $\pi(F_i) = E_{i+1,i}$ and $\pi(K_i) = I + (q-1)E_{ii}$, where E_{ij} are the matrices $(E_{ij})_{rs} = \delta_{ir}\delta_{js}$.

We now turn to the description of the center $\mathcal{Z}$ of the quantum $\mathrm{U}_q(\mathfrak{gl}_n)$. Let $\mathrm{U}_q(\mathfrak{gl}_n)^-$, $\mathrm{U}_q(\mathfrak{gl}_n)^+$ and $\mathrm{U}_q(\mathfrak{gl}_n)^0$ the subalgebras of $\mathrm{U}_q(\mathfrak{gl}_n)$ generated by $F_{i'}$ ($i' \in I'$), $E_{i'}$ ($i' \in I'$) and $K_i^{\pm}$ ($i \in I$) respectively. Any element $z \in \mathcal{Z}$ can be written as

$$z = z^{(0)} + \sum_s u_s^{(-)} u_s^{(0)} u_s^{(+)},$$

where $z^{(0)}, u_s^{(0)} \in \mathrm{U}_q(\mathfrak{gl}_n)^0$, $u_s^{(+)} \in \mathrm{U}_q(\mathfrak{gl}_n)^+$ and $u_s^{(-)} \in \mathrm{U}_q(\mathfrak{gl}_n)^-$. The quantum Harish-Chandra homomorphism is an algebra homomorphism $\varphi : \mathcal{Z} \longrightarrow \mathrm{U}_q(\mathfrak{gl}_n)^0$ such that $z \mapsto z^{(0)}$.

The dot action of the Weyl group W of $\mathfrak{gl}_n$ on $\mathrm{U}_q(\mathfrak{gl}_n)^0$ is given by permutations of the elements $q^{-i}K_i$ ($i \in \mathbf{I}$). Define

$$L_i = q^{(\varepsilon_i, 2\rho)}(K_i)^2, \quad i \in \mathbf{I}. \tag{2.1}$$

Then L_i are permuted by the Weyl group. Let $\mathrm{U}_q(\mathfrak{gl}_n)^0_{ev}$ be the subalgebra spanned by the elements $\Pi_{i=1}^n L_i^{l_i}$ for $l_i \in \mathbb{Z}$. We denote by $(U_q(\mathfrak{gl}_n)^0_{ev})^W$ the W-invariant subalgebra of $\mathrm{U}_q(\mathfrak{gl}_n)^0_{ev}$. By using the quantum Harish-Chandra isomorphism for $U_q(\mathfrak{sl}_n)$ (see, e.g., [2, 7]), one can prove the following result.

Lemma 2.1. *The Harish-Chandra homomorphism is an algebra isomorphism between the center $\mathcal{Z}$ of $\mathrm{U}_q(\mathfrak{gl}_n)$ and the subalgebra of $\mathrm{U}_q(\mathfrak{gl}_n)^0$ generated by the elements of $(U_q(\mathfrak{gl}_n)^0_{ev})^W$ together with c, where*

$$c = K_1^{-1}K_2^{-1}\cdots K_n^{-1}. \tag{2.2}$$

Note that c is obviously W-invariant, and $c^{\pm 2} \in (U_q(\mathfrak{gl}_n)^0_{ev})^W$. An equivalent description of this lemma can be found in [1]. In this paper, a set of generators for $\mathcal{Z}$ were also given, which are different from the quantum Casimirs operators of [8, 9].

2.2. **Quantum Casimir operators of** $\mathrm{U}_q(\mathfrak{gl}_n)$.

The quantum Casimir operators of $\mathrm{U}_q(\mathfrak{gl}_n)$ are the main objects for study in this paper, which we now briefly describe. As is well known, in the quantum group setting, we have neither a good quantum analogue of tensors nor a procedure for "contracting tensors" (however, see [11, 12]). Thus it is much harder to explicitly construct central elements for quantum groups. The construction of [8, 9] was actually quite involved: it had to invoke the theory of [10] and also made use the universal R-matrix of $\mathrm{U}_q(\mathfrak{gl}_n)$. Thus for the sake of completeness and also clearness we briefly explain the construction.

It is well-known that $\mathrm{U}_q(\mathfrak{gl}_n)$ is a quasi-triangular Hopf algebra, i.e., there exists an invertible element $R \in \mathrm{U}_q(\mathfrak{gl}_n) \otimes \mathrm{U}_q(\mathfrak{gl}_n)$ which is called the universal R-matrix of

$\mathrm{U}_q(\mathfrak{gl}_n)$, such that

$$\begin{gathered} R\Delta(x)=\Delta'(x)R, \quad \forall x\in \mathrm{U}_q(\mathfrak{gl}_n),\\ (\Delta\otimes id)R=R_{13}R_{23}, \quad (id\otimes\Delta)R=R_{13}R_{12}, \end{gathered}$$

where Δ' is the opposite co-multiplication. Explicitly, $\Delta'=T\circ\Delta$ where $T:\mathrm{U}_q(\mathfrak{gl}_n)\otimes \mathrm{U}_q(\mathfrak{gl}_n)\to \mathrm{U}_q(\mathfrak{gl}_n)\otimes\mathrm{U}_q(\mathfrak{gl}_n)$ is the linear map defined for any $x,y\in\mathrm{U}_q(\mathfrak{gl}_n)$ by $T(x\otimes y)=y\otimes x$. The R-matrix satisfies the celebrated Yang-Baxter equation

$$R_{12}R_{13}R_{23}=R_{23}R_{13}R_{12}.$$

Denote $R^T=T(R)$. Then $R^TR\Delta(x)=\Delta(x)R^TR$, $\forall\, x\in\mathrm{U}_q(\mathfrak{gl}_n)$

For any $\alpha\in\mathbb{Z}$, Let $\mathbb{Z}_{\geqslant\alpha}:=\{m\,|\,m\in\mathbb{Z},\, m\geqslant\alpha\}$. The following quantum Casimir operators for $\mathrm{U}_q(\mathfrak{gl}_n)$ were constructed in [8, 9]:

$$C_{n,k}=\mathrm{Tr}_\pi\left(1\otimes q^{\pi(2h\rho)}\left(\frac{\Gamma-1\otimes 1}{q-q^{-1}}\right)^k\right),\ \ k\in\mathbb{Z}_{\geqslant 0},$$

where Tr_π represents the trace taken over π, and

$$\Gamma=(id\otimes\pi)R^TR,\quad q^{\pi(2h\rho)}=\prod_{i=1}^n q^{(n-2i+1)E_{i,i}}.$$

Let L_Λ be a finite dimensional irreducible $\mathrm{U}_q(\mathfrak{gl}_n)$-module with highest weight $\Lambda\in\mathcal{H}^*$. Then each $C_{n,k}$ acts on L_Λ by a scalar, which is given by the following formula [4]:

$$\begin{aligned}\chi_\Lambda(C_{n,k}) \;=\; & (q-q^{-1})^{-k}\sum_{i=1}^n\left(q^{(\varepsilon_i,2\Lambda+2\rho+\varepsilon_i)-C(\Lambda_0)}-1\right)^k\\ & \times q^{C(\Lambda_0)-(\varepsilon_i,\varepsilon_i)}\prod_{j\neq i}^n\frac{q^{(\varepsilon_i,2\Lambda+2\rho+\varepsilon_i)}-q^{(\varepsilon_j,2\Lambda+2\rho-\varepsilon_j)}}{q^{(\varepsilon_i,2\Lambda+2\rho+\varepsilon_i)}-q^{(\varepsilon_j,2\Lambda+2\rho+\varepsilon_j)}},\end{aligned}$$

where $\Lambda_0=\varepsilon_1$.

Since $\varphi(z)$ belongs to $\mathrm{U}_q(\mathfrak{gl}_n)^0$ for every $z\in\mathcal{Z}$, the right hand side of the above formula must be a polynomial in $q^{(\varepsilon_i,2\Lambda+2\rho)}$. Note that when applying L_i to the highest weight vector v_Λ of L_Λ, we have $L_iv_\Lambda=q^{2(\varepsilon_i,\Lambda+\rho)}v_\Lambda$. Then it follows that

$$C^0_{n,k}:=\varphi(C_{n,k})=\sum_{i=1}^n\Big(\frac{q^{1-n}L_i-1}{q-q^{-1}}\Big)^k\prod_{j\neq i}^n\frac{qL_i-q^{-1}L_j}{L_i-L_j}. \tag{2.3}$$

3. ANALYSIS OF $G_{n,k}$

Let us analyze the formula for $C^0_{n,k}$ to put it into a form which will be readily usable for the proof of Theorem 4.1. Denote

$$G_{n,k}=\sum_{i=1}^n {L_i}^kP_{n,i}\ \text{ where }\ P_{n,i}=\prod_{j\neq i}^n\frac{qL_i-q^{-1}L_j}{L_i-L_j}.$$

Then for any $n \in \mathbb{Z}_{\geqslant 2}$ and $k \in \mathbb{Z}_{\geqslant 1}$, one can rewrite $C^0_{n,k}$ as

$$C^0_{n,k} = \frac{1}{(q^{-1}-q)^k} \sum_{j=0}^{k} \left(\begin{array}{c} k \\ j \end{array} \right) (-q^{1-n})^j G_{n,j}.$$

We want to prove that $G_{n,k}\,(k = 0,1,\cdots,n)$ are polynomials in L_i.

Lemma 3.1. *For any $n \geqslant 2$, the following two identities hold:*

$$G_{n,0} \;=\; q^{n-1} + q^{n-3} + \cdots + q^{3-n} + q^{1-n}, \tag{3.1}$$

$$G_{n,1} \;=\; q^{n-1}(L_1 + L_2 + \cdots + L_n). \tag{3.2}$$

Proof For $i \neq n, n-1$, one has

$$\begin{aligned} P_{n,i} \;&=\; P_{n-2,i} \cdot \frac{qL_i - q^{-1}L_{n-1}}{L_i - L_{n-1}} \cdot \frac{qL_i - q^{-1}L_n}{L_i - L_n} \\ &=\; P_{n-2,i} \cdot \frac{qL_i - q^{-1}L_{n-1}}{L_i - L_{n-1}} \cdot \frac{qL_i - q^{-1}L_n}{L_{n-1} - L_n} - P_{n-2,i} \cdot \frac{qL_i - q^{-1}L_n}{L_i - L_n} \cdot \frac{qL_i - q^{-1}L_{n-1}}{L_{n-1} - L_n}. \end{aligned}$$

For any $m, n \in \mathbb{Z}_{\geqslant 2}$ and $i \in \mathbb{Z}_{\geqslant 0}$, denote $P_{m,i,n} = P_{m,i}\frac{qL_i - q^{-1}L_n}{L_i - L_n}$. Then $P_{n,i}$ can be rewritten as

$$P_{n,i} \;=\; P_{n-2,i,n-1} \cdot \frac{qL_i - q^{-1}L_n}{L_{n-1} - L_n} - P_{n-2,i,n} \cdot \frac{qL_i - q^{-1}L_{n-1}}{L_{n-1} - L_n}. \tag{3.3}$$

For convenience, denote $G_{n,k,\widehat{j}} = \sum\limits_{j \neq i=1}^{n} {L_i}^k P_{n,i}$. Then we can rewrite $G_{n,0}$ as follows:

$$\begin{aligned} G_{n,0} \;&=\; \sum_{i=1}^{n-2} P_{n-2,i,n-1} \cdot \frac{qL_i - q^{-1}L_n}{L_{n-1} - L_n} + P_{n,n-1} - \sum_{i=1}^{n-2} P_{n-2,i,n} \cdot \frac{qL_i - q^{-1}L_{n-1}}{L_{n-1} - L_n} + P_{n,n} \\ &=\; \frac{qG_{n-1,1} - q^{-1}L_n G_{n-1,0}}{L_{n-1} - L_n} - \frac{qG_{n,1,\widehat{n-1}} - q^{-1}L_{n-1}G_{n,0,\widehat{n-1}}}{L_{n-1} - L_n}. \qquad (3.4) \end{aligned}$$

Meanwhile, for any $n \in \mathbb{Z}_{\geqslant 2}$ and $k \in \mathbb{Z}_{\geqslant 1}$, we have the following computations:

$$\begin{aligned} G_{n,k} - L_n G_{n,k-1} \;&=\; \sum_{i=1}^{n} L_i^{k-1}(L_i - L_n) P_{n,i} \\ &=\; \sum_{i=1}^{n-1} L_i^{k-1} P_{n-1,i}(qL_i - q^{-1}L_n) \\ &=\; qG_{n-1,k} - q^{-1}L_n G_{n-1,k-1}. \end{aligned}$$

Thus

$$G_{n,k} \;=\; L_n G_{n,k-1} + qG_{n-1,k} - q^{-1}L_n G_{n-1,k-1}, \;\; \forall\, n \in \mathbb{Z}_{\geqslant 2}, k \in \mathbb{Z}_{\geqslant 1}. \tag{3.5}$$

It is easy to see that both (3.1) and (3.2) hold for the cases $n = 2$ and $n = 3$. Conbining the identities (3.4) and (3.5), using induction on n in (3.1) and (3.2), one can get the formulae for $G_{n,0}$ and $G_{n,1}$. Hence we complete the proof of this lemma. □

We introduce a generating function $S_n(t) = \sum_{k=0}^{\infty} t^k G_{n,k}$. Then

$$S_n(t) - G_{n,0} - tL_nS_n(t) = qS_{n-1}(t) - q^{-1}tL_nS_{n-1}(t) - qG_{n-1,0},$$

which implies

$$\begin{aligned} S_n(t) &= \frac{q - q^{-1}tL_n}{1 - tL_n} S_{n-1}(t) + \frac{G_{n,0} - qG_{n-1,0}}{1 - tL_n} \\ &= \frac{q - q^{-1}}{1 - tL_n} S_{n-1}(t) + q^{-1}S_{n-1}(t) + \frac{q^{-(n-1)}}{1 - tL_n}. \end{aligned}$$

Thus

$$G_{n,k} = qG_{n-1,k} + (q - q^{-1}) \sum_{i=0}^{k-1} L_n^{k-i} G_{n-1,i} + q^{-(n-1)} L_n^k. \tag{3.6}$$

Note that it is by no means obvious from formula (2.3) itself that its right hand side is a polynomial in L_i's. To put our mind at peace, we observe the following result.

Lemma 3.2. *For any $n \in \mathbb{Z}_{\geqslant 2}$ and $k \in \mathbb{Z}_{\geqslant 0}$, $G_{n,k}$ is a polynomial in L_i's.*

Proof. This lemma follows from Lemma 3.1 and the identities (3.5) and (3.6). □

Form the formulae proved in Lemma 3.1, we see that if L_i is replaced by e^{ε_i} for $i = 1, 2, \cdots, n$, then the coefficient of q^{n-1} in $G_{n,1}$ corresponds to the character of the basic irreducible representation of $\mathrm{U}_q(\mathfrak{gl}_n)$. A natural problem is to understand all the $G_{n,k}$ in similar terms, and we address this problem now.

Let us introduce a set of elements of λ_k^i in $\sum_{i=1}^n \mathbb{Z}_+ \varepsilon_i$, which we write in terms of their coordinates relative to the basis ε_i. We let $\lambda_k^1 = (k, 0, 0, \cdots, 0)$, $\lambda_k^2 = (k-1, 1, 0, \cdots, 0)$, $\cdots$, $\lambda_k^k = (1, \cdots, 1, 0, \cdots, 0)$ in the case $k < n$. We also set $\lambda_k^1 = (k, 0, 0, \cdots, 0)$, $\lambda_k^2 =$

$(k-1,1,0,\cdots,0)$, $\cdots$, $\lambda_k^k=(k-n+1,1,\cdots,1)$ in the case $k\geqslant n$. Note that these weights respectively correspond to the following Young diagrams:



Denote by $ChL_{\lambda_k^i}$ the character of the irreducible $\mathfrak{gl}_n$-representation with highest weight λ^i. Then

$$ChL_{\lambda_k^i}=\frac{\sum\limits_{w\in W} sign(w)e^{w((k-i+1)\varepsilon_1+\varepsilon_2+\cdots+\varepsilon_i+\widetilde{\rho})}}{\prod\limits_{i<j}(e^{\varepsilon_i}-e^{\varepsilon_j})},$$

where $\widetilde{\rho}=\rho+\frac{1}{2}(n-1)\sum_{i=1}^n\varepsilon_i$. The formula is valid when $k-i\geq 0$. When $i>k$, the right hand vanishes identically.

Lemma 3.3. *Let $Ch_{n,k}$ be the expression obtained from $G_{n,k}$ by replacing L_i by e^{ε_i} for all $i=1,2,\cdots,n$. Then*

$$\sum_{i=1}^n(-1)^{i-1}q^{n-2i+1}ChL_{\lambda_k^i}=Ch_{n,k},\quad \forall\, k=1,2,\cdots, \tag{3.7}$$

where $ChL_{\lambda_k^i}=0$ if $k<i$.

Proof Denote $W_{n-1}=\{w\in W\mid w(\varepsilon_1)=\varepsilon_1\}$. Then $W/W_{n-1}=\{[w]\mid w\in W\mid w(\varepsilon_1)=\varepsilon_j,\ \forall\, j\neq 1\}$. Denote $\widetilde{\rho}_{n-1}=\widetilde{\rho}-(n-1)\varepsilon_1$. In the case when $k\geqslant n$, the left side of

(3.7) can be rewritten as follows:

$$\begin{aligned}
&\frac{q^{n-1}\sum\limits_{w\in W} sign(w)w\big(e^{k\varepsilon_1+\widetilde{\rho}}\prod\limits_{i=2}^{n}(1-q^{-2}e^{-(\varepsilon_1-\varepsilon_i)})\big)}{\prod\limits_{i<j}(e^{\varepsilon_i}-e^{\varepsilon_j})}\\
&=q^{n-1}\sum_{w\in W} w\frac{e^{k\varepsilon_1+\widetilde{\rho}}\prod\limits_{i=2}^{n}(1-q^{-2}e^{-(\varepsilon_1-\varepsilon_i)})}{\prod\limits_{i<j}(e^{\varepsilon_i}-e^{\varepsilon_j})}\\
&=q^{n-1}\sum_{[\sigma]\in W/W_{n-1}}\sum_{w'\in W_{n-1}}\sigma\Big(w'\big(\frac{e^{\widetilde{\rho}_{n-1}}}{\prod\limits_{1<i<j}(e^{\varepsilon_i}-e^{\varepsilon_j})}\big)e^{(k+n-1)\varepsilon_1}\prod_{j=2}^{n}\frac{1-q^{-2}e^{-(\varepsilon_1-\varepsilon_j)}}{e^{\varepsilon_1}-e^{\varepsilon_j}}\Big).
\end{aligned}$$

According to the fact $ChL_0=1$, we have the following identity

$$\sum_{[w']\in W/W_{n-1}} w'\frac{e^{\widetilde{\rho}_{n-1}}}{\prod\limits_{i<j}(e^{\varepsilon_i}-e^{\varepsilon_j})}=1.$$

Hence the left side of (3.7) can be further simplified as follows:

$$\begin{aligned}
&q^{n-1}\sum_{\sigma\in W/W_{n-1}}\sigma(e^{(k+n-1)\varepsilon_1}\prod_{j=2}^{n}\frac{1-q^{-2}e^{-(\varepsilon_1-\varepsilon_j)}}{e^{\varepsilon_1}-e^{\varepsilon_j}})\\
&=\sum_{[\sigma]\in W/W_{n-1}}\sigma(e^{k\varepsilon_1}\prod_{j=2}^{n}\frac{qe^{\varepsilon_1}-q^{-1}e^{\varepsilon_j}}{e^{\varepsilon_1}-e^{\varepsilon_j}}),
\end{aligned}$$

which is just the right side of (3.7), i.e. $G_{n,k}$ after replacing e^{ε_i} by L_i for $i=1,2,\cdots,n$. Thus the lemma in the case $k\geqslant n$ follows.

In the case $1\leqslant k<n$, the proof can be proceed similarly by recalling the fact $ChL_{\lambda_k^i}=0$ for $i>k$. We omit the details. $\square$

We have the following immediate consequence of the above lemma.

Corollary 3.1. *Set $\Gamma_{k,i}=0$ if $i>k$. For $i\leqslant k$, let $\Gamma_{k,i}$ be obtained by replacing each e^{ε_i} in $ChL_{\lambda_k^i}$ by $L_i\in \mathrm{U}_q(\mathfrak{gl}_n)^0$. Then*

$$G_{n,k} \quad = \quad \sum_{i=1}^{n}(-1)^{i-1}q^{n-2i+1}\Gamma_{k,i}.$$

Note that the nonzero $\Gamma_{k,i}$ are symmetric polynomials in L_1, L_2, $\cdots$, L_n.

Let $\mathcal{G}_{n,k}$ denote the rational function in q obtained by replacing e^{ε_i} by $q^{2(\varepsilon_i,\Lambda+\rho)}$ for all $i=1,2,\cdots,n$ on the left side of (3.7). Then we obtain a new formula for the eigenvalues of the quantum Casimir operators of $\mathrm{U}_q(\mathfrak{gl}_n)$ in the irreducible representation with highest weight Λ.

Corollary 3.2.

$$\chi_\Lambda(C_{n,k}) = (q-q^{-1})^{-k}\sum_{l=0}^{k}(-1)^{k-l}\left(\begin{array}{c} k \\ l \end{array}\right)(q^{1-n})^l\mathcal{G}_{n,l}.$$

The advantage of this formula is that every term in $\mathcal{G}_{n,l}$ has a representation theoretical interpretation in terms of the general linear Lie algebra.

4. THE MAIN RESULT

With the preparations in the previous sections, we can now prove the following theorem, which is the main result of the paper.

Theorem 4.1. *The center* $\mathcal{Z}$ *of* $\mathrm{U}_q(\mathfrak{gl}_n)$ *is generated by* c *and the quantum Casimir operators* $C_{n,1}, \cdots, C_{n,n}$.

In order to prove the theorem, we need some basic results on symmetric polynomials [6], which we recall here. The complete homogeneous symmetric polynomial of degree k in n variables x_1, x_2, $\cdots$, x_n, written h_k for $k = 0, 1, 2, \cdots$, is the sum of all monomials of total degree k in the variables. Formally,

$$h_k(x_1, x_2, \cdots, x_n) = \sum_{1\leqslant i_1\leqslant i_2\leqslant\cdots\leqslant i_k\leqslant n} x_{i_1}x_{i_2}\cdots x_{i_k}.$$

It is well known that the set of complete homogeneous symmetric polynomials

$$h_1(x_1, x_2, \cdots, x_n), h_2(x_1, x_2, \cdots, x_n), \cdots, h_n(x_1, x_2, \cdots, x_n)$$

generate the ring of symmetric polynomials in the n variables x_1, x_2, $\cdots$, x_n.

Proof of Theorem 4.1. Denote $\mathbb{K}[L_1, L_2, \cdots, L_n]^W$ the algebra of symmetric polynomials in the polynomial ring $\mathbb{K}[L_1, L_2, \cdots, L_n]$. Given any element in $(U_q(\mathfrak{gl}_n)^0_{ev})^W$, we can always express it in terms of elements of $\mathbb{K}[L_1, L_2, \cdots, L_n]^W$ and c algebraically. Note that $c^{-2} \in \mathbb{K}[L_1, L_2, \cdots, L_n]^W$. Therefore, in order to prove Theorem 4.1, it suffices to show that $G_{n,k}\,(k = 1, \cdots, n)$ generate $\mathbb{K}[L_1, L_2, \cdots, L_n]^W$.

Note that $\Gamma_{k,1}$ in $G_{n,k}$ is a complete symmetric polynomial in L_1, $\ldots$, L_n. Thus $\Gamma_{1,1}$, $\Gamma_{2,1}$, $\ldots$, $\Gamma_{n,1}$ are a set of generators of the ring $\mathbb{K}[L_1, L_2, \cdots, L_n]^W$ of symmetric polynomial.

Now $\Gamma_{1,1}$ is equal to $q^{1-n}G_{n,1}$, and we can easily express $\Gamma_{2,1}$ in terms of $G_{n,1}$ and $G_{n,2}$. Inductively we can show that $\Gamma_{k,1}$ can always be expressed in terms of $G_{n,1}$, $G_{n,2}$, $\cdots$, $G_{n,k}$. Thus $G_{n,1}$, $G_{n,2}$, $\cdots$, $G_{n,n}$ are also a set of generators of the symmetric polynomial ring $\mathbb{K}[L_1, L_2, \cdots, L_n]^W$. Since the elements of $\mathbb{K}[L_1, L_2, \cdots, L_n]^W$ and c together generate $\varphi(\mathcal{Z})$, we complete the proof of the theorem. □

Acknowledgements This work is supported by the Australian Research Council. Financial assistance from the National Science Foundation of China (Grant No. 10926166) is also acknowledged.

REFERENCES

[1] M.J Hopkins, A.I. Molve, A q-analogue of the centralizer construction and skew representations of the quantum Affine algebra, *Symmetry, Intergrability and Geometry: methods and applications* **2** (2006), 092-1–092-29.

[2] J.C. Jantzen, Lectures on quantum groups, Graduate Studies in Mathematics 6, Amer. Math. Soc., Providence, RI, 1996.

[3] M. Kashiwara, Crystallizing the q-analogue of universal enveloping algebras, Commun. Math. Phys. 133 (1990), 249–260.

[4] J. R. Links, R. B. Zhang, Eigenvalues of Casimir invariants of $U_q(\mathrm{gl}(m|n))$, *J. Math. Phys.* **34** (1993), 6016–6024.

[5] G. Lusztig, Canonical bases arising from quantized enveloping algebras, J. Amer. Math. Soc. 3 (1990), no. 2, 447498

[6] I. G. Macdonald, Symmetirc functions and Hall polynomials, second ed. Oxford: Clarendon Press, (1995).

[7] T. Tanisaki, Harish-Chandra isomorphism for quantum groups, *Commun. Math. Phys.* **127** (1990), 555–571.

[8] Gould, M. D.; Zhang, R. B.; Bracken, A. J. Generalized Gel?fand invariants and characteristic identities for quantum groups. J. Math. Phys. 32 (1991), no. 9, 2298–2303.

[9] R. B. Zhang, M. D. Gould, A. J. Bracken, Generalized Gel'fand invariants of quantum groups, *J. Phys. A*, **24** (1991), 937–943.

[10] Zhang, R. B.; Gould, M. D.; Bracken, A. J. Quantum group invariants and link polynomials. Comm. Math. Phys. 137 (1991), no. 1, 13–27.

[11] Zhang, R. B.; Gould, M. D.; Bracken, A. J. From representations of the braid group to solutions of the Yang-Baxter equation. Nuclear Phys. B 354 (1991), no. 2-3, 625–652.

[12] Rittenberg, V.; Scheunert, M. Tensor operators for quantum groups and applications. J. Math. Phys. 33 (1992), no. 2, 436–445.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA